

APPENDIX

NSUN3 and ABH1 modify the wobble position of mt-tRNA^{Met} to expand codon recognition in mitochondrial translation

Sara Haag, Katherine E. Sloan, Namit Ranjan, Ahmed S. Warda, Jens Kretschmer, Charlotte Blessing, Benedikt Hübner, Jan Seikowski, Sven Dennerlein, Peter Rehling, Marina V. Rodnina, Claudia Höbartner and Markus T. Bohnsack

TABLE OF CONTENTS

Table S1

Table S2

Supplementary Methods

Table S1: Oligonucleotides used in this study

Name	Sequence (5'-3')
NSUN3_qPCR_fwd	TGGGTCTGTTTGGAAATCCTATT
NSUN3_qPCR_rev	TGCACCACCTTAAATCATTGTTAC
ABH1_qPCR_fwd	CACCATTCTGCTGTGCCCTA
ABH1_qPCR_rev	CAAGCTGCCTACCCTCAGAC
GAPDH_qPCR_fwd	CTGGCGTCTTCACCACCATGG
GAPDH_qPCR_rev	CATCAGCCACAGTTTCCCGG
mt-tRNA ^{Met} _RT	TAATAATACAAAAAATATAACCAAC
anti-mt-tRNA ^{Met}	GGGAAGGGTATAACCAACATTTTCGGGGTATGGGCCCGATAGCTTAT TTAGCTGACC
anti-mt-tRNA ^{Glu}	CACGGACTACAACCACGACCAATGATATGAAAAACCATCGTTGTATT CAACTACAAG
anti-mt-tRNA ^{Pro}	GAGAAAAAGTCTTTAACTCCACCATTAGCACCCAAAGCTAAGATTCTA ATTTAACTATTCTC
anti-tRNA ^{Met} _i	GGTTTCGATCCATCGaCCTCTGGGTTATGGGCCAGCAGCCTTCCGC TGCG
anti-tRNA ^{Met} _e	GAGGCTTGAActCAGGaCCTTCAGATTATGAGACTGACGCGCTGCCA GCTG

Table S2: Antibodies used in this study

Name	Source/Reference
ABH1	Abcam (ab128895)
TIM44	Proteintech (13859-1-AP)
TIM23	Dennerlein et al, 2015
TOM70	Dennerlein et al, 2015
FLAG	Sigma Aldrich (F3165)
Tubulin	Sigma Aldrich (T6199)

Table S3: siRNAs used in this study

Name	Sequence (5'-3')
siABH1_1	UGACCAGAAUAGCGAAGUA
siABH1_3	GUGGUGACAUCAUGAUAUU
siNSUN3_1	GGACUUGCUCACGACUU
siNSUN3_2	GCAAAGUUGUGUUGGAUCA
siNT	UCGUAAGUAAGCGCAACCC

Supplementary Methods

Crosslinking and analysis of cDNA (CRAC)

Crosslinking and analysis of cDNA experiments were performed as previously described (Bohnsack et al, 2012). In brief, HEK293 cells expressing NSUN3-HisPrcFlag, NSUN3-C265A-HisPrcFLAG, ABH1-HisPrcFLAG or the HisPrcFlag tag alone were induced using 1 µg/mL tetracycline for 24 h. UV crosslinking was carried out using a Stratalinker (Stratagene) as previously described (Sloan et al, 2015). For chemical crosslinking cells were incubated in growth media complemented with 4 µM 5-azacytidine (Sigma) as previously described (Haag et al, 2015). Cells were harvested in buffer containing 50 mM Tris/HCl pH 7.6, 150 mM NaCl, 0.1% NP-40, 5 mM β-mercaptoethanol and protease inhibitors, and lysed by sonication. Protein-RNA complexes were affinity purified using anti-FLAG magnetic beads (Sigma) before elution using PreScission Protease. A partial RNase digest (1 U RNaseH1 (Agilent) for 30 sec at 37°C) was performed and complexes were purified under denaturing conditions (6 M guanidium-HCl) on Ni-NTA (Qiagen). Co-purified RNAs were radiolabelled using T4 PNK (Thermo Scientific) and ³²P-γ-ATP (Perkin Elmer) and RNA-protein complexes were separated by NuPAGE gel electrophoresis then transferred to a nitrocellulose membrane. Radioactive signals were detected by exposure to an X-ray film. For Illumina sequencing, 3' and 5' adapters were ligated to the co-precipitated RNA and a cDNA library was prepared by reverse transcription of the RNA using Superscript III Reverse Transcriptase (Thermo) and amplification using primers containing randomised five-nucleotide sequence to enable PCR templates to be distinguished. Before mapping Flexbar was used to remove 3'-adapter sequences and bases with a phred quality score <13 (95% base call accuracy) and identical sequence reads were collapsed using python scripts. Bowtie2 was used to map the remaining reads on the human ensembl genome version GRCh 37.75 with an 18-nucleotide cut-off.

Expression and purification of recombinant proteins

Expression of His₁₄-MBP-NSUN3 or His₁₄-MBP-ABH1 (and mutants thereof) in *Escherichia coli* (DE3) Rosetta pLysS cells was induced by addition of 0.5 mM IPTG for 16 h at 18°C. Cells were harvested and lysed by sonication in a buffer containing 30 mM KPi pH 7.0, 300 mM KCl, 10% (v/v) glycerol, 10 mM imidazole, 0.1 mM dithiothreitol (DTT) and protease inhibitors (complete mini, Roche). The lysate was cleared by centrifugation and His-tagged proteins were retrieved on cComplete His-Tag purification resin (Roche). After washing steps with lysis buffer, proteins were eluted in a buffer composed of 30 mM KPi pH 7.0, 300 mM KCl, 10% glycerol, 200 mM imidazole and 0.1 mM DTT. In the case of NUSN3, the eluate was diluted five-fold in a buffer containing 20 mM KPi pH 7.0, 200 mM KCl, 10% (v/v) glycerol, 2 mM beta-mercaptoethanol and then incubated with amylose resin. Elution of bound proteins was achieved using the dilution buffer supplement with 10 mM maltose. All proteins were dialysed against a buffer containing 30 mM KPi pH 7.0, 100 mM KCl, 50% glycerol, 1 mM DTT, and 0.1 mM EDTA for storage.

For MTIF2-His₆ or TUFM-His₆, expression in BL21 Codon Plus cells was induced by addition of 1 mM IPTG for 4 h at 37°C before cells were harvested. Cells were resuspended in a buffer containing 50 mM Tris-HCl pH 8.0, 60 mM NH₄Cl, 7 mM MgCl₂, 300 mM KCl, 10 mM imidazol, 10% (v/v) glycerol, 5 mM beta-mercaptoethanol, 50 µM GDP and lysed using a “French press”. After pelleting cell debris by centrifugation, the cleared lysate was incubated with Ni-NTA (Qiagen). Washing steps were performed with lysis buffer before elution in buffer composed of 50 mM Tris-HCl pH 8.0, 60 mM NH₄Cl, 7 mM MgCl₂, 300 mM KCl, 250 mM imidazol, 10% (v/v) glycerol, 5 mM beta-mercaptoethanol, 50 µM GDP. Proteins were further purified on a Sephadex 75 column equilibrated in 50 mM Tris-HCl pH 7.5, 70 mM NH₄Cl, 7 mM MgCl₂, 30 mM KCl and fractions containing the MTIF2 or TUFM were pooled and concentrated.

Preparation of [¹⁴C]Met-tRNA^{Met} and ribosome binding assays

Aminoacylation of mitochondrial tRNAs was carried out in 160 µL reaction volume containing 50 mM HEPES pH 7.5, 70 mM NH₄Cl, 30 mM KCl, 11 mM MgCl₂, 3 mM ATP, 2 mM β-mercaptoethanol, 5 A₂₆₀ units tRNAs (unmodified, m⁵C, f⁵C, hm⁵C), 20 µM [¹⁴C]Methionine, and 15% (v/v) of *Escherichia coli* methionine tRNA synthetase (MetRS). After incubation for 30 min at 37°C, aminoacylated tRNA was extracted with phenol and precipitated with 2% (w/v) KAc pH 5.0 and cold ethanol.

Ribosome binding experiments were carried out in buffer TAKM₇ (50 mM Tris-HCl pH 7.5, 50 mM NH₄Cl, 50 mM KCl, 7 mM MgCl₂, 1 mM DTT) if not stated otherwise.

Ribosomes, EF-Tu, and fMet-tRNA^{fMet} from *E. coli* were prepared as described (Rodnina and Wintermeyer, 1995; Gromadski et al, 2006; Milon et al, 2007). Preparation of POST complex was carried out as previously described (Milon et al, 2007; Belardinelli et al, 2016). Ribosomes (4 μ M) were incubated with a two-fold excess of mRNA (GGCAAGGAGGUAAAUA AUG UUU AUA GUU AC, or GGCAAGGAGGUAAAUA AUG UUU AUG GUU AC (IBA Göttingen); codon occupying the A-site is underlined) in the presence of 6 μ M initiation factors IF1, IF2, IF3, 6 μ M f³H]Met-tRNA^{fMet}, and 1 mM GTP in TAKM₇ for 30 min at 37°C. Ternary complexes of EF-Tu–GTP–Phe-tRNA^{Phe} were prepared by incubating EF-Tu with 1 mM GTP, 3 mM phosphophenolpyruvate and 0.1 mg/ml pyruvate kinase for 15 min at 37 °C prior to the addition of Phe-tRNA^{Phe}. The POST complex was formed by mixing initiation complex with ternary complex for 1 min at RT and with EF-G (0.1 μ M) for an additional 1 min at room temperature. The resulting POST complexes were purified by centrifugation through a 1.1 M sucrose cushion in buffer A. For mitochondrial [¹⁴C]Met-tRNA^{Met} binding to the A site, ternary complexes of TUFM–GTP–[¹⁴C]Met-tRNA^{Met} were prepared by incubating TUFM with 1 mM GTP, 3 mM phosphophenolpyruvate, 0.1 mg/ml pyruvate kinase and [¹⁴C]Met-tRNA^{Met} prepared from unmodified, m⁵C, hm⁵C or f⁵C tRNAs for 20 min at 30°C. The ternary complex was added to the POST complex and incubated for 10 min at room temperature. The amount of f³H]Met-Phe-[¹⁴C]Met bound to ribosomes was determined by nitrocellulose filtration.

For mitochondrial [¹⁴C]Met-tRNA^{Met} binding to the P-site, 1 μ M ribosomes were incubated with a two-fold excess of mRNA (GGCAAGGAGGUAAAUA AUG AAA UUU GUU AC, or GGCAAGGAGGUAAAUA AUA AAA UUU GUU AC, or GGCAAGGAGGUAAAUA AUU UUU UAA AGU UAC (IBA Göttingen); codon occupying the P-site is underlined) in the presence of 2 μ M human mitochondrial initiation factor 2 (MTIF2), 0.15 μ M [¹⁴C]Met-tRNA^{Met} prepared from unmodified, m⁵C, hm⁵C or f⁵C tRNAs and 1 mM GTP in TAKM₇ for 10 min at 30°C. The amount of [¹⁴C]Met bound to ribosomes was determined by nitrocellulose filtration and scintillation counting.

Supplementary References

Belardinelli R, Sharma H, Caliskan N, Cunha CE, Peske F, Wintermeyer W, Rodnina MV (2016) Choreography of molecular movements during ribosome progression along mRNA. *Nat Struct Mol Biol.* 23: 342-348

Bohnsack MT, Tollervey D, Granneman S (2012) Identification of RNA helicase target sites by UV cross-linking and analysis of cDNA. *Methods Enzymol* 511: 275-288

Dennerlein S, Oeljeklaus S, Jans D, Hellwig C, Bareth B, Jakobs S, Deckers M, Warscheid B, Rehling P (2015) MITRAC7 Acts as a COX1-Specific Chaperone and Reveals a Checkpoint during Cytochrome c Oxidase Assembly. *Cell Rep* 12: 1644-1655

Gromadski KB, Daviter T, Rodnina MV (2006) A uniform response to mismatches in codon-anticodon complexes ensures ribosomal fidelity. *Mol Cell* 21: 369-377

Haag S, Warda AS, Kretschmer J, Günningmann MA, Höbartner C, Bohnsack MT (2015) NSUN6 is a human RNA methyltransferase that catalyzes formation of m5C72 in specific tRNAs. *RNA* 21: 1532-1543

Milon P, Konevega AL, Peske F, Fabbretti A, Gualerzi CO, Rodnina MV (2007) Transient kinetics, fluorescence, and FRET in studies of initiation of translation in bacteria. *Methods Enzymol* 430: 1-30

Rodnina MV, Wintermeyer W (1995) GTP consumption of elongation factor Tu during translation of heteropolymeric mRNAs. *Proc Natl Acad Sci USA* 92: 1945-1949

Sloan KE, Leisegang MS, Doebele C, Ramírez AS, Simm S, Safferthal C, Kretschmer J, Schorge T, Markoutsas S, Haag S, Karas M, Ebersberger I, Schleiff E, Watkins NJ, Bohnsack MT (2015) The association of late-acting snoRNPs with human pre-ribosomal complexes requires the RNA helicase DDX21. *Nucleic Acids Res* 43: 553-564

Expanded View Figures




Figure EV1. The cytoplasmic tRNA Met-i and Met-e do not represent substrates of NSUN3 *in vivo*.

- A, B The UV or 5-AzaC cross-linking and analysis of cDNA (CRAC) experiments with NSUN3-HisPrcFLAG or FLAG control cells were performed as described for Fig 2. (A) The percentages of the Illumina sequence reads mapped to individual classes of RNA are given graphically for each sample. Abbreviations: tRNA, transfer RNA; snRNA, small nuclear RNA; snoRNA, small nucleolar RNA; rRNA, ribosomal RNA; mtRNA, mitochondrial-encoded RNA; miscRNA, miscellaneous RNA; miRNA, microRNA; lncRNA, long non-coding RNA. (B) The relative distribution of cytoplasmic tRNA sequence reads obtained from the CRAC experiments is shown. Only tRNAs that were represented by more than 5% of all cytoplasmic tRNA reads are labelled.
- C *In vitro* methylation reactions were performed using recombinant His₁₄-MBP-NSUN3 (NSUN3) or the catalytically inactive mutant His₁₄-MBP-NSUN3-C265A (C265A), [³H-methyl]-labelled S-adenosylmethionine as a methyl group donor and *in vitro*-transcribed mitochondrial mt-tRNA^{Met}, cytoplasmic tRNA₁^{Met} and tRNA_e^{Met}. The RNA was then separated on a denaturing polyacrylamide gel, stained with ethidium bromide (EtBr) to indicate inputs and exposed to an X-ray film to analyse methylation (³H-Me).
- D 5-AzaC cross-linking was performed and RNA-associated with wild-type NSUN3, the catalytic NSUN3 mutant (C265A) or the FLAG tag alone was isolated as described in (A). The RNA was isolated from the purified protein–RNA complexes and analysis by Northern blot using probes against the mt-tRNA^{Met}, mt-tRNA₁^{Met} and mt-tRNA_e^{Met}. Inputs are shown on the left and eluates on the right. The mt-tRNA^{Met} panel is identical to that shown in Fig 2G.
- E The nucleotide sequences of the anticodon stem loops of mt-tRNA^{Met} (left) and tRNA₁^{Met} (right) are shown.




Figure EV2. The mitochondrial pool of ABH1 is localised in the matrix.

To analyse submitochondrial localisation of ABH1, human mitochondria were isolated and either left untreated, swollen in hypotonic buffer (Mitoplasts) or disrupted by sonication (Sonic.) before treatment with different amounts of proteinase K (PK) where indicated, followed by SDS-PAGE and Western blotting using antibodies against human TIM44, TIM23, TOM70 or FLAG-tagged NSUN3. Note that TIM44 extrudes into the matrix, while a major domain of TIM23 localises to the intermembrane space and TOM70 is largely exposed on the mitochondrial surface. The asterisk indicates a cross-reaction of the TOM70 antibody. The panels of TIM44, TIM23 and TOM70 are identical with those shown in Fig 1B.




Figure EV3. RNAi-mediated depletion of NSUN3 or ABH1 leads to reduced cell growth.

HeLa cells that had been transfected with siRNAs against NSUN3 (siNSUN3_1 or siNSUN3_2), ABH1 (siABH1_1 or siABH1_2) or non-target siRNAs (siNTs) were harvested and counted. Cell numbers from three experiments were normalised to the non-target control, and results are given graphically as mean ± SD.